

The Wu-Yang Ambiguity Revisited*Daniel Z. Freedman¹ and Ramzi R. Khuri²

*Theory Division
CERN
CH-1211 Geneva 23
Switzerland*

ABSTRACT

Several examples are given of continuous families of $SU(2)$ vector potentials $A_i^a(x)$ in 3 space dimensions which generate the same magnetic field $B^{ai}(x)$ (with $\det B \neq 0$). These Wu-Yang families are obtained from the Einstein equation $R_{ij} = -2G_{ij}$ derived recently via a local map of the gauge field system into a spatial geometry with 2-tensor $G_{ij} = B^a{}_i B^a{}_j \det B$ and connection Γ^i_{jk} with torsion defined from gauge covariant derivatives of B .

Submitted to: *Physics Letters B*

CERN-TH.7187/94
CTP#2282
hep-th 9403031
March 1994

* This work is supported in part by funds provided by the National Science Foundation (NSF) Grant PHY/9206867 and the U. S. Department of Energy (DOE) under contract #DE-AC02-76ER03069.

¹ Permanent address: Department of Mathematics and Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139-4307, USA.

² Supported by a World Laboratory Fellowship.

The Wu-Yang ambiguity [1] is the phenomenon that two or more gauge inequivalent non-abelian potentials $A_i^a(x)$ generate the same field strength $F_{ij}^a(x)$. This was widely discussed in the literature [2-14] of the 1970's. Although the original example is 3-dimensional, it was mainly the 4-dimensional case which was of past interest. Many examples of a discrete ambiguity have been exhibited, specifically two potentials A and $\bar{A}$ giving the same F . The few examples of a continuous ambiguity [7-14] were degenerate in some way: for example, they were effectively 2-dimensional.

In this communication we study the Wu-Yang ambiguity of $SU(2)$ gauge field configurations in 3-spatial dimensions, and we obtain several examples of continuous families of potentials which generate the same magnetic field

$$B^{ai} = \epsilon^{ijk} \left[\partial_j A_k^a + \frac{1}{2} \epsilon^{abc} A_j^b A_k^c \right]. \quad (1)$$

In 3 dimensions there is no “algebraic obstruction” to an ambiguity. Given two potentials, A and $\bar{A}$, we find by subtraction of their Bianchi identities that their difference $\Delta A = A - \bar{A}$ satisfies

$$\epsilon^{ijk} \epsilon^{abc} B^{ki} \Delta A_j^c = 0 \quad (2)$$

which is an underdetermined linear system — 3 equations for 9 unknowns — for any B -field configuration. In four dimensions one gets a “square” homogeneous linear system — 12 equations for 12 unknowns — which has a non-trivial solution only if the field strength satisfies an algebraic condition corresponding to vanishing determinant [5,6,8]. In any case, (2) is not sufficient to demonstrate that (1), viewed as a partial differential equation for A_j^b given B^{ai} , has multiple solutions, and it is this which we wish to explore here, largely by means of examples.

The 3-dimensional case is relevant for the Hamiltonian form of gauge field dynamics in $3 + 1$ dimensions and especially for attempts [15], [16] to transform from A_i^a to B^{ai} as the basic field variables. In fact the work here emerged from further study of the recent proposal [16] to do this as an intermediate step to replace the A, B system by a set of gauge-invariant spatial geometric variables, namely a metric G_{ij} and connection Γ_{ij}^k with torsion. It turns out that the information we find on the Wu-Yang ambiguity invalidates the form of Hamiltonian dynamics proposed in [16]. But the geometry described there is valid, and it is through the geometrical equations that our Wu-Yang information is obtained. See [17] for an earlier and similar geometrical treatment of the equations of motion of 2+1 dimensional $SU(2)$ Yang-Mills theory.

1. We begin with our first example. Consider the smooth, algebraically non-singular (*i.e.* $\det B \neq 0$) magnetic field $B^{ai} = \delta^{ai}$, in Euclidean space with Cartesian coordinates x, y, z . It is easy to show explicitly that, for any real parameter β with $|\beta| > 1$, the 1-parameter family of potentials

$$A_i^a = \begin{pmatrix} \beta \pm \sqrt{\beta^2 - 1} \cos(z/\beta) & \pm \sqrt{\beta^2 - 1} \sin(z/\beta) & 0 \\ \pm \sqrt{\beta^2 - 1} \sin(z/\beta) & \beta \mp \sqrt{\beta^2 - 1} \cos(z/\beta) & 0 \\ 0 & 0 & 1/\beta \end{pmatrix} \quad (3)$$

all reproduce the same B^{ai} . Gauge inequivalence is demonstrated by the fact that the invariants $B^{aj}D_iB^{ak}$ depend on β and z .

- a. Note the “nonlinear wave” character of these potentials — the amplitude and wavelength of the trigonometric terms are correlated.
- b. The particular magnetic field $B^{ai} = \delta^{ai}$ is invariant under rotations and translations of the configuration space $\mathbb{R}^3$ (the spatial rotations must be combined with a suitably chosen $\text{SO}(3)$ gauge transformation, constant in this case, as is well known). Since eq. (1) is also covariant, each such isometry which does not leave A_i^a invariant produces another Wu-Yang related potential. In this way one can extend the potentials displayed in (3) to a 4-parameter family, in which the wave has an arbitrary phase, $z \rightarrow z - z_0$ and direction $(0, 0, 1) \rightarrow \hat{k}$.
- c. Brown and Weisberger [11] are often credited with the proof that a uniform non-abelian field strength such as our B^{ai} also requires a uniform vector potential, so there is an apparent conflict with the decidedly non-uniform potentials displayed in (3). However Brown and Weisberger define a uniform field by the strong requirement that all gauge invariant quantities, e.g. $B^{ai}D_jB^{ak}$, are constant, so there is no conflict with their work.
- d. Equation (1) is covariant under diffeomorphisms $y^\alpha(x^i)$ of $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ under which A and B transform as follows:

$$\begin{aligned} A'^a_\alpha(y) &= \frac{\partial x^i}{\partial y^\alpha} A^a_i(x) \\ B'^{a\alpha}(y) &= \left| \frac{\partial x}{\partial y} \right| \frac{\partial y^\alpha}{\partial x^i} B^{ai}(x) \end{aligned} \tag{4}$$

Thus examples of the Wu-Yang ambiguity automatically extend to entire orbits of the diffeomorphism group. The magnetic energy

$$E = \frac{1}{2} \int d^3x \delta_{ij} B^{ai}(x) B^{aj}(x) \tag{5}$$

involves the fixed Cartesian metric δ_{ij} , and is not invariant under diffeomorphisms. Below we will show that one can find a diffeomorphism under which the field $B^{ai} = \delta^{ai}$ which has infinite energy, transforms to a configuration $B'^{a\alpha}(y)$ which falls sufficiently fast as $y^\alpha \rightarrow \infty$ that it has finite energy.

2. Let us now review the spatial geometry of [16] which is the main tool used in this work. Let $B^a_i(x)$ denote the matrix inverse of the magnetic field of $\text{SU}(2)$ gauge theory, and $\det B = \det B^{ai}$, which is gauge invariant. Then $G_{ij}(x) = B^a_i(x)B^a_j(x)/\det B$ is a gauge invariant symmetric tensor (under diffeomorphisms)*. The following geometry, which obviously uses the fact that the gauge group $\text{SU}(2)$ is also the tangent space group of a 3-manifold, emerged from the physical aim of studying the action of the electric field on gauge invariant state functionals $\psi[G]$.

* One sees that G_{ij} is a positive definite (negative definite) tensor, when $\det B > 0$ ($\det B < 0$) We discuss the positive case here because only that is used below. See [16] for a complete treatment.

The quantity $b_i^a = |\det B|^{\frac{1}{2}} B_i^a$ is essentially a frame for G_{ij} . One may apply a Yang-Mills covariant derivative and define a quantity Γ_{ij}^k as follows:

$$D_i b_j^a = \Gamma_{ij}^k b_k^a. \quad (6)$$

It then follows immediately that Γ_{ij}^k transforms as a connection under diffeomorphisms and that it is well defined in terms of A_i^a and B^{ai} for all non-singular magnetic fields, $\det B \neq 0$. If we multiply (6) by b_k^a and symmetrize in j, k , we easily derive

$$\partial_i G_{jk} - \Gamma_{ij}^l G_{lk} - \Gamma_{ik}^l G_{jl} = 0. \quad (7)$$

This means that Γ is a metric compatible connection for G , and can be written as

$$\Gamma_{ij}^k = \dot{\Gamma}_{ij}^k(G) - K_{ij}{}^k \quad (8)$$

where $\dot{\Gamma}$ is the Christoffel connection and K is the contortion tensor, which is antisymmetric in the last pair of indices, $K_{ijk} = -K_{ikj}$.

Next one studies the anti-symmetrized Yang-Mills derivative of (6)

$$[D_i, D_j] b_k^a = b_l^a \partial_i \Gamma_{jk}^l + \Gamma_{jk}^l \partial_i b_l^a - (i \leftrightarrow j) \quad (9)$$

One uses (6) again to find that the curvature tensor of Γ appears. On the left one uses the gauge field Ricci identity and a standard formula for the inverse of a 3×3 matrix. The result is

$$R_{kij}^l = \delta_j^l G_{ik} - \delta_i^l G_{jk}, \quad (10)$$

the statement that the spatial geometry associated with the gauge fields A, B is maximally symmetric. However, in 3 dimensions, with or without torsion, the Riemann tensor R_{kij}^l is fully determined by its Ricci contraction $R_{kj} = \delta_l^i R_{kij}^l$, so that no information is lost by restricting consideration to the contracted form of (8), namely

$$R_{ij}(\Gamma) = -2G_{ij} \quad (11)$$

which defines an Einstein geometry with torsion. One may show [16] using the second Bianchi identity of curvatures with torsion, that an integrability condition for (11) is that the contortion tensor is traceless, $K_{kj}{}^k = 0$, and that this is also a direct requirement of the gauge field Bianchi identity, $D_i B^{ai} = 0$, applied to the definition (6) of Γ .

The discussion above defines the forward map from Yang-Mills fields A and B , always related by (1), to geometric variables G and Γ defined by explicit local formulas above. The gauge field Ricci and Bianchi identities then imply that G and Γ are related by the Einstein condition (11) with traceless contortion. The fundamental reason for the Einstein geometry is that the magnetic field is simultaneously the curvature (1) of the gauge connection A and also essentially the frame of the spatial geometry.

One may also ask about the inverse map from tensor $G_{ij}(x)$ and connection $\Gamma_{ij}^k(x)$ on $\mathbb{R}^3$ to gauge fields. Suppose that a frame b_i^a , with $\det b > 0$, is constructed for G_{ij} by any standard method, then (6) can be written out as

$$\partial_i b_j^a - \Gamma_{ij}^k b_k^a + \epsilon^{abc} A_i^b b_j^c = 0. \quad (12)$$

This is just the “dreibein postulate” with A essentially the spin connection, and one can solve for A , obtaining [16]

$$A_i^a = -\frac{1}{2} \epsilon^{abc} b^{bj} (\partial_i b_j^c - \Gamma_{ij}^k b_k^c) \quad (13)$$

while the magnetic field is defined from the inverse frame by

$$B^{ai}(x) = |\det G_{jk}|^{\frac{1}{2}} b^{ai}. \quad (14)$$

Thus given a frame one obtains the magnetic field from (14), while both b and Γ are required to define the potential via (13). Since the frame is unique up to a local $SO(3)$ rotation, these maps define A and B uniquely up to an $SU(2)$ gauge transformation. Furthermore, A and B defined in this way satisfy the gauge theory relation (1) if Γ and G satisfy (11).

Thus the gauge theory Wu-Yang ambiguity will appear whenever the Einstein equation (11) viewed as a partial differential equation for K , given G , has multiple solutions. To investigate this it is useful to use the representation [16]

$$K^i_{jk} = \epsilon_{jkn} S^{ni} \frac{1}{|\det G|^{1/2}} \quad (15)$$

which automatically satisfies the antisymmetry and tracelessness requirements if S^{ni} is a symmetric tensor. When (11) is expanded out using (8) and (15), one finds that the Einstein equation is equivalent to

$$\frac{\epsilon^{jkl}}{|\det G|^{1/2}} \dot{\nabla}_k S_{li} - \left(S_k^j S_i^k - S_k^k S_i^j \right) = \dot{R}_i^j + 2\delta_i^j. \quad (16)$$

In (16) $\dot{\nabla}_k$ indicates a spatial covariant derivative with Christoffel connection $\dot{\Gamma}$ and $\dot{R}_{ij}$ is the conventional symmetric Ricci tensor. The $\epsilon \dot{\nabla} S$ term is non-symmetric, so that (16) comprises 9 equations for the 6 components of S_{ij} . However it was shown explicitly in [16] that there is a Bianchi identity which imposes 3 constraints on the 9 equations, so there is no reason to think that (16) is an overdetermined system. From (8), (13) and (15), A can be expressed in terms of S as

$$A_i^a = -\frac{1}{2} \epsilon^{abc} b^{bj} \dot{\nabla}_i b_j^c - b^{ak} S_{ki}. \quad (17)$$

Our approach to the Wu-Yang ambiguity is to take an input metric $G_{ij}(x)$ and study the solutions of (16) for the torsion $S_{ij}(x)$. It is not clear why this should be a simpler method than to study directly whether (1) has multiple solutions for A , given B . Perhaps it is because an equation for the 6 components of S_{ij} is simpler to handle than an equation for

the 9 components of A_i^a , but it may just be an historical accident that has led us to approach the Wu-Yang ambiguity via the spatial geometry of [16].

Before beginning to study applications of (16), it is perhaps useful to note that (6) indicates that Γ is completely determined by first covariant derivatives $D_i B^{aj}$ of the magnetic field. It then follows from properties of the inverse map discussed above that there is no Wu-Yang ambiguity for the potential A_i^a , if we require that both B and DB are preserved^{**}. In 4 dimensional $SU(2)$ gauge theory [2,3], the field strength and its first two covariant derivatives determine the potential locally uniquely.

The Wu-Yang ambiguity indicates that the potential $A_i^a(x)$ contains gauge invariant information beyond that in the magnetic field $B^{ai}(x)$. Therefore the change of field variable $A_i^a \rightarrow B^{ai}$ which was the basis of the version of gauge invariant Hamiltonian dynamics presented in [16] is invalid. The discrete 2:1 ambiguity envisaged there could be handled, but it is probably impossible to deal with a continuous ambiguity without serious revision of the proposal. In the 2+1 Lagrangian theory the equation of motion $D^\mu F_{\mu\nu} = 0$ provides additional information, presumably enough to fix the Wu-Yang ambiguity, which then should not be a difficulty for [17].

3. It is the tensor $G_{ij}(x) = \delta_{ij}$ that corresponds to the magnetic field $B^{ai} = \delta^{ai}$, and it can be seen without difficulties that the torsion solutions of (14) are related in this simple case to the potentials of (3) by $A_i^a(z) = -S_{ai}(z)$. Note that at $\beta = 1$, the solution (3) reduces to $S_{ij} = \delta_{ij}$, and was already noted in [16]. We found the family of solutions (3) by first linearizing about $S_{ij} = \delta_{ij}$ and using Fourier analysis to find linearized modes of wave number $k^2 = 1$. This led us to investigate the single variable ansatz $S_{ij}(z)$ which reduces (16) to a non linear system of ordinary differential equations. Some fiddling then led to (3), which is unique within this ansatz (except for translation $z \rightarrow z - z_0$). One can show that the only spherically symmetric solutions of (16) for input $G_{ij} = \delta_{ij}$ are the solutions $S_{ij} = \pm\delta_{ij}$ recognized in [16]. There is a heuristic argument that the potentials displayed in (3) together with those obtained from them by translation and rotation are the only potentials for the field $B^{ai} = \delta^{ai}$ which continuously limit to potentials $\bar{A}_i^a = \delta_i^a$ with $\beta = 1$ in (3). The reason is that one can show using the Fourier transform that the set of linear perturbations about $\bar{A}_i^a$ obtained in the $\beta \rightarrow 1$ limit of our Wu-Yang family are complete.

We now discuss the use of diffeomorphisms in relation to the energy of field configurations exhibiting the Wu-Yang ambiguity. We proceed in several steps.

- a. Given an initial configuration $B^{ai}(x)$ in Cartesian coordinates x^m on $\mathbb{R}^3$, we transform to spherical coordinates $y^i = (r, \theta, \varphi)$. Both the spacetime metric δ_{ij} and the magnetic field $B^{ai}(x)$ transform to

$$\begin{aligned} \delta_{ij} &\rightarrow g_{ij}(y) = \frac{\partial x^m}{\partial y^i} \frac{\partial x^n}{\partial y^j} \delta_{mn}, \\ B^{ai} &\rightarrow B'^{ai}(y) = \left| \frac{\partial x}{\partial y} \right| \frac{\partial y^i}{\partial x^m} B^{am}(x), \end{aligned} \tag{18}$$

^{**} Actually it is sufficient to require that $D_{[i} b_{j]}^a$ is preserved, because this determines the torsion tensor from (6).

and the energy (5) can be rewritten as

$$E = \frac{1}{2} \int d^3y \frac{1}{\sqrt{g(y)}} g_{ij}(y) B'^{ai} B'^{aj} = \frac{1}{2} \int d^3y \frac{1}{\sqrt{g(y)}} g_{ij}(y) G'^{ij}(y) \det G'. \quad (19)$$

For the specific case of spherical coordinates and field $B^{ai} = \delta^{ai}$ this is equal to the divergent expression

$$E = \frac{3}{2} \int dr d\theta d\varphi r^2 \sin \theta, \quad (20)$$

which could have been obtained by naive change of variables in (5).

- b. We now make the diffeomorphism $y^i \rightarrow z^i$ of the vector potential and the magnetic field by changing the radial coordinate

$$\begin{aligned} z^1 &= R(r), \\ z^2 &= \theta = y^2, \\ z^3 &= \varphi = y^3, \\ \tilde{B}^{ai} &= \left| \frac{\partial y}{\partial z} \right| \frac{\partial z^i}{\partial y^m} B'^{am}(y), \\ \tilde{G}^{ij} &= \frac{\partial z^i}{\partial y^m} \frac{\partial z^j}{\partial y^n} G'^{mn}(y). \end{aligned} \quad (21)$$

In the specific case $B^{ai} = \delta^{ai}$, G'^{mn} is the standard diagonal inverse metric for (r, θ, φ) , and $\tilde{G}^{ij}$ is also diagonal and given by

$$\begin{aligned} \tilde{G}^{RR} &= R'^2, \\ \tilde{G}^{\theta\theta} &= \frac{1}{r^2}, \\ \tilde{G}^{\varphi\varphi} &= \frac{1}{r^2 \sin^2 \theta}, \end{aligned} \quad (22)$$

- c. The coordinates $z^\alpha = (R, \theta, \varphi)$ are new spherical coordinates on $\mathbb{R}^3$, so the energy of the $\tilde{B}^{ai}(z)$ field configuration is

$$\tilde{E} = \frac{1}{2} \int d^3z \frac{1}{\sqrt{g(z)}} g_{ij}(z) \tilde{G}^{ij}(z) \det \tilde{G}(z), \quad (23)$$

with g_{ij} the standard metric for (R, θ, φ) . For the example $B^{ai} = \delta^{ai}$, one can use (18-23) and obtain

$$\tilde{E} = \frac{1}{2} \int dR d\theta d\varphi \frac{r^4 \sin \theta}{R^2} \left(1 + \frac{2}{R'^2} \right). \quad (24)$$

The constraints that $\tilde{B}^{ai}(z)$ is a finite energy field configuration on $\mathbb{R}^3$ are that $R(r)$ map $(0, \infty) \rightarrow (0, \infty)$ monotonically such that r^4/R^2 falls faster than $1/R$ and no other singularities appear. It is not difficult to show that $R = r\sqrt{1+r^8}$ satisfies these requirements. It is certainly not unique.

4. The next set of Wu-Yang examples we discuss emerge from the 3-dimensional hyperbolic metrics

$$ds^2 = \frac{1}{c^2 z^2} (dx^2 + dy^2 + dz^2) \quad (25)$$

for which $R_{ij} = -2c^2 G_{ij}$. One may anticipate that the case $c^2 = 1$ is especially simple because the right side of (16) vanishes. It turns out that one can also make the $\epsilon \dot{\nabla} S$ and SS terms vanish separately. There is no integrability constraint when $\dot{\nabla}_j$ is applied to the former condition, while the second condition implies that S_{ij} is a rank 1 dyadic matrix. With this structure in view one can easily find that within the two variable ansatz $S_{ij}(z, x)$, there is the family of solutions

$$S_{ij}(z, x) = \delta_{i1} \delta_{j1} \frac{1}{z} h(x) \quad (26)$$

which involves an arbitrary function of the variable x . The solution can be rotated by an angle θ in the x, y plane to obtain

$$\begin{aligned} S_{ij} &= \frac{1}{z} h(x \cos \theta + y \sin \theta) V_i V_j, \\ V_i &= (\cos \theta, \sin \theta, 0). \end{aligned} \quad (27)$$

We have not studied the application of the full $SO(2, 1) \times SO(2, 1)$ isometry group of the metric (25), but more solutions seem likely. In this frame the magnetic field is given simply by $B^{ai} = \delta^{ai}/z^2$ while the gauge potential corresponding to (26) is obtained from equation (17):

$$A_1^1 = -h(x), \quad A_2^1 = -A_1^2 = 1/z, \quad (28)$$

with the rest of the components vanishing. In this frame, however, the magnetic field B^{ai} is singular on the plane $z = 0$, so we transform our configuration to a frame in which both the magnetic field and gauge potential are manifestly regular over all of $\mathbb{R}^3$. Transforming to the coordinates

$$\begin{aligned} u^1 &= \frac{4x}{x^2 + y^2 + (z+1)^2}, \\ u^2 &= \frac{4y}{x^2 + y^2 + (z+1)^2}, \\ u^3 &= \frac{2(x^2 + y^2 + z^2 - 1)}{x^2 + y^2 + (z+1)^2}, \end{aligned} \quad (29)$$

the metric can be re-written in the form

$$ds^2 = \frac{1}{(1 - r^2/4)^2} ((du^1)^2 + (du^2)^2 + (du^3)^2), \quad (30)$$

where $r^2 = (u^1)^2 + (u^2)^2 + (u^3)^2 < 4$, which is another standard metric on a 3-hyperboloid and can be analytically continued to a metric on a 3-sphere. The inverse transformation is

then given by

$$\begin{aligned} x &= \frac{u^1}{(u^1/2)^2 + (u^2/2)^2 + (u^3/2 - 1)^2}, \\ y &= \frac{u^2}{(u^1/2)^2 + (u^2/2)^2 + (u^3/2 - 1)^2}, \\ z &= \frac{1 - ((u^1/2)^2 + (u^2/2)^2 + (u^3/2)^2)}{(u^1/2)^2 + (u^2/2)^2 + (u^3/2 - 1)^2}. \end{aligned} \quad (31)$$

We make a further transformation to the coordinates

$$v^i = u^i / (1 - r^2/4) \quad (32)$$

so that the entire $\mathbb{R}^3$ is covered. The inverse transformation is given by

$$u^i = \frac{2(\sqrt{1+v^2} - 1)}{v^2} v^i \quad (33)$$

so that the metric can be written in the form

$$ds^2 = \frac{dv^2}{1+v^2} + dv^2 d\Omega^2 = dv^i dv^i - \left(\frac{1}{1+v^2} \right) v^i dv^i v^j dv^j. \quad (34)$$

From the last form, one can define a frame and use (14) to obtain the following regular configuration for the B -field:

$$B^{ai} = \frac{\delta^{ai}}{\sqrt{1+v^2}} + \left(1 - \frac{1}{\sqrt{1+v^2}} \right) \frac{v^a v^i}{v^2}, \quad (35)$$

in which v^i are regarded as Cartesian coordinates. A dyadic solution for S in the (xyz) frame then transforms to a dyadic solution in the $(v^1 v^2 v^3)$ frame according to

$$S'_{ij}(v^k) = S_{mn}(x^p) \frac{\partial x^m}{\partial v^i} \frac{\partial x^n}{\partial v^j}. \quad (36)$$

In particular, the solution $S_{11}(x, z) = h(x)/z$ transforms into

$$S'_{ij}(v^k) = \frac{1}{z(v^k)} h(x(v^k)) V_{(i)} V_{(j)}, \quad (37)$$

where $V_{(i)} = \partial x / \partial v^i$ and where $x(v^k)$ and $z(v^k)$ are easily given from (31) and (33). Using equation (17) we compute the corresponding set of potentials A_i^a in the v^k -frame all of which yield the magnetic field of (35):

$$A_i^a = -\epsilon^{aik} \left(\frac{\sqrt{1+v^2} - 1}{v^2} \right) v^k - S'_{ai} - \left(\frac{\sqrt{1+v^2} - 1}{v^2} \right) v^a v^k S'_{ki}. \quad (38)$$

When $c^2 \neq 1$ the full nonlinear equations are very difficult to handle, so we restrict ourselves to a perturbative expansion about the symmetric solution $\bar{S}_{ij} = \sqrt{1 - c^2} G_{ij}$ of (16) by setting $S_{ij} = \bar{S}_{ij} + \hat{\Sigma}_{ij}$. The perturbation $\hat{\Sigma}_{ij}$ satisfies the linear equation

$$(cz)\epsilon^{jkl}\dot{\nabla}_k\hat{\Sigma}_{li} + \sqrt{1 - c^2}(\hat{\Sigma}_{ji} + \hat{\Sigma}_{kk}\delta_{ji}) = 0. \quad (39)$$

(the placement of the j index reflects the removal of the conformal factor.) The 9 equations for the 6 components of $\hat{\Sigma}$ cannot all be independent, and the fact that we find a consistent solution below is a practical test of the exact Bianchi identity [16] satisfied by (16). Note that the ij contraction of (39) immediately tells us that the trace $\hat{\Sigma}_{kk} = 0$.

Because of the x -translation symmetry of the metric (25) we look for a solution of the form $\hat{\Sigma}_{ij}(z, x) = \Sigma_{ij}(z, k)e^{ikx}$. The 9 equations of (39) can be manipulated to obtain a second order differential equation for the component Σ_{23}

$$\left(z^2 \frac{d^2}{dz^2} + z \frac{d}{dz} - k^2 z^2 + \frac{1 - c^2}{c^2}\right) \Sigma_{23} = 0, \quad (40)$$

while the other components are related to Σ_{23} by

$$\begin{aligned} \Sigma_{33} &= \frac{-ikc}{\sqrt{1 - c^2}} z \Sigma_{23}, \\ \Sigma_{13} &= \frac{c}{\sqrt{1 - c^2}} z \frac{d}{dz} \Sigma_{23}, \\ \Sigma_{12} &= \frac{i}{k} \left(\frac{d}{dz} - \frac{1}{z} \right) \Sigma_{23}, \\ \Sigma_{11} &= \frac{-ic}{k\sqrt{1 - c^2}} z \frac{d^2}{dz^2} \Sigma_{23}, \\ \Sigma_{22} &= -(\Sigma_{11} + \Sigma_{33}). \end{aligned} \quad (41)$$

Note that (40) is the differential equation for Bessel functions of imaginary argument ikz and index $p = \sqrt{(c^2 - 1)/c^2}$ which is also imaginary when $c^2 < 1$ and the symmetric torsion $\bar{S}_{ij}$ is real.

Note that the wave number k of the linear perturbation is not restricted in contradistinction to the flat metric where, as can be seen from the small amplitude limit $\beta \rightarrow 1$ in (3), the wave number $k = 1$ is required. This means that the general real superposition

$$\int dk \varphi(k) \Sigma_{ij}(kz) e^{ikx} + \text{c.c.} \quad (42)$$

is also a solution, so we have the freedom of an arbitrary function at the linear level. We expect that (42) can be used as the “input” to the system of differential equations determining second and higher order perturbative solutions of (16), and that the functional freedom of $\varphi(k)$ remains. Thus the qualitative picture of the torsion solutions for the hyperbolic metrics for all values of c is that they contain an arbitrary function of a single variable and the additional

parametric freedom obtained from isometries. The case $c = 1$ is special only because exact solutions can be easily obtained.

5. The final set of Wu-Yang ambiguities we discuss arise from the $2 + 1$ product metrics of the form

$$ds^2 = F(x, y)(dx^2 + dy^2) + dz^2, \quad (43)$$

and for which $\dot{R}_{ij} = -\delta_{ij}\nabla^2 \ln F$ and $\dot{R}_{i3} = \dot{R}_{33} = 0$ for $i, j = 1, 2$. Then provided F satisfies the differential equation

$$\nabla^2 \ln F = c^2 F + \frac{m^2}{F}, \quad (44)$$

where $c^2 < 1$ and $m > 0$, a solution for S_{ij} is given by

$$S_{ij} = \begin{pmatrix} \beta F \pm m \cos(z/\beta) & \pm m \sin(z/\beta) & 0 \\ \pm m \sin(z/\beta) & \beta F \mp m \cos(z/\beta) & 0 \\ 0 & 0 & 1/\beta \end{pmatrix} \quad (45)$$

where $\beta = \sqrt{1 - c^2}$. For $m \neq 0$, this solution represents a continuous 1-parameter family of Wu-Yang ambiguities obtained by translations in z .

For $m = 0$ the metric (43) reduces to the constant curvature case, the wave disappears, and the torsion becomes

$$\tilde{S}_{11} = \tilde{S}_{22} = \beta F, \quad \tilde{S}_{33} = 1/\beta. \quad (46)$$

The manifestly symmetric form of the product metric is

$$ds^2 = \frac{1}{c^2 y^2} (dx^2 + dy^2) + dz^2, \quad (47)$$

and we now study the perturbative expansion about the symmetric $\tilde{S}_{ij}$ of (46) by setting $S_{ij} = \tilde{S}_{ij} + \tilde{\Sigma}_{ij}$. Because of the z -translation symmetry of the metric (47) we look for a solution of the form $\tilde{\Sigma}_{ij}(y, z) = \Sigma'_{ij}(y, k)e^{ikz}$. The 9 equations in the linearized approximation can be manipulated to obtain a second order differential equation for the component Σ'_{23}

$$\left(\frac{d^2}{dy^2} + \frac{2}{y} \frac{d}{dy} + \frac{\alpha(a, k)}{y^2} \right) \Sigma'_{23} = 0, \quad (48)$$

where $\alpha(\beta, k) = 2\beta^2(\beta^2 - k^2)/(1 - \beta^4)$ while the other components are simply related to Σ'_{23} . The solution of (48) is given simply by the power behaviour $\Sigma'_{23} = Ay^t$, where A is an arbitrary constant and $t = -1/2 \pm \sqrt{1/4 - \alpha(\beta, k)}$. For $\tilde{\Sigma}_{ij}(y, x) = \Sigma''_{ij}(y, k)e^{ikx}$, Σ''_{ij} is simply expressed in terms of Bessel functions of imaginary argument iky and index $m = \sqrt{1/4 - 2\beta^2/(1 - \beta^4)}$, which is either real or imaginary depending on the value of β .

Again the wave number k of the linear perturbation is not restricted and we expect that this functional freedom persists in higher order perturbative solutions.

6. What we have exhibited in this paper are several examples of a continuous Wu-Yang ambiguity for $SU(2)$ gauge fields in 3 dimensions and a new technique, namely the

Einstein space condition (14) for obtaining such field configurations. It is intriguing to ask about the systematics of the ambiguity; namely what properties of the B -field determine the degree of ambiguity in the associated potentials A . Our examples provide at least a limited view of this systematics. Certainly an ambiguity is generated whenever there is a symmetry transformation of B which acts nontrivially on A , but this is not enough to explain the free parameter β in (3), nor the arbitrary functions such as $F(x)$ in the example (26-38) or in the various linear solutions we have presented. Gauge field topology does not seem to be the issue here for two reasons. First of all the ambiguity can be exhibited in any compact subset of the configuration space $\mathbb{R}^3$. Second, if we are given in some gauge a Wu-Yang family with suitable behavior at spatial infinity one can apply a gauge transformation to change the topological class at will. Of course one does expect that, except for singularities of the map (1), the degree of ambiguity in A will not change as the parameters of B are smoothly varied. Our examples appear to be consistent with this requirement, although the discrete ambiguity found when $\dot{R}_{ij} = 0$ must be understood as a singular limit of the case of non-zero curvature. It is interesting that the Riemannian curvature $\dot{R}_{ij}$ of the metric G_{ij} obtained from B plays a role both in the ease of obtaining solutions for A and in the qualitative nature of the ambiguity.

Acknowledgements

We wish to thank L. Alvarez-Gaumé, M. Bauer, S. Deser, C. Imbimbo, J. Liu and E. Verlinde for helpful discussions.

References

- [1] T.T. Wu and C.N. Yang, *Phys. Rev.* **D12** (1975) 3845.
- [2] C.H. Gu and C.N. Yang, *Sci. Sin.* **18** (1975) 484; *Sci. Sin.* **20** (1977) 47.
- [3] C.L. Shen, *Fudan Journal* (Natural Science) **2** (1976) 61.
- [4] S. Deser and F. Wilczek, *Phys. Lett.* **65B** (1976) 391.
- [5] R. Roskies, *Phys. Rev.* **D15** (1977) 1731.
- [6] M. Calvo, *Phys. Rev.* **D15** (1977) 1733.
- [7] S. Coleman, *Phys. Lett.* **70B** (1977) 59.
- [8] M.B. Halpern, *Phys. Rev.* **D16** (1977) 1798; *Nucl. Phys.* **B139** (1978) 477.
- [9] S. Solomon, *Nucl. Phys.* **B147** (1979) 174.
- [10] C.G. Bollini, J.J. Giambiagi and J. Tiomno, *Phys. Lett.* **83B** (1979) 185.
- [11] L.S. Brown and W.I. Weisberger, *Nucl. Phys.* **B157** (1979) 285.
- [12] S. Deser and W. Drechsler, *Phys. Lett.* **86B** (1979) 189.
- [13] M. Mostow, *Commun. Math. Phys.* **78** (1980) 137.
- [14] F.A. Doria, *Commun. Math. Phys.* **79** (1981) 435; *J. Math. Phys.* **22** (1981) 2943.
- [15] A.M. Badalyan, *Sov. J. Nucl. Phys.* **38** (1983) 464.
- [16] D.Z. Freedman, P.E. Haagensen, K. Johnson and J.I. Latorre, CERN-TH.7010/93, CTP #2238, hep-th 9309045 (September 1993).
- [17] F.A. Lunev, *Phys. Lett.* **B295** (1992) 99.